

Submentale Laserassistierte Liposuktion: Klinische Erfahrungen und Histologische Ergebnisse

Submental Laser-Assisted Liposuction:

Clinical Experience and Histological Findings

Alberto Goldman, MD, Porto Alegre, RS, Brazil

Schlüsselwörter

Laserlipolyse, Laserassistierte Liposuktion, Lipoplastik, Liposkulptur

Key words

Laserlipolysis, laser-assisted liposuction, lipoplasty, liposculpture

Zusammenfassung

Die submentale Liposuktion ist ein exzellentes Verfahren zur Verjüngung des alternden Halses, indem der zervikom mentale Winkel reduziert und die Konturen der Kiefergrenze hervorgehoben werden. Die Liposuktion ist zur Zeit eine der am häufigsten durchgeführten cosmeto-chirurgischen Prozeduren. Fortschritte in der Kanüलगestaltung, neue Konzepte, und verschiedene technische Optionen haben die Ergebnisse am Patienten verbessert. Bedenken gibt es weiterhin bezüglich des Gesamtvolumens des Aspirats, sowie der möglichen klinischen und hämodynamischen Risiken beim Patienten, also auch der postoperativen Genesung und des Endergebnisses. Liposuktion des Halses und der Backen ist klinisch gesehen, eines der lohnendsten Verfahren in der ästhetischen Chirurgie. In der hier präsentierten Studie, beschreibt der Autor seine Erfahrungen in der laserassistierte Liposuktion des Halses und der Backen sowie die Auswirkungen auf das Fett- und umliegende Gewebe.

In diesem Verfahren wird die submental lokalisierte Lipodystrophie mit einem Neodymium-, Yttrium-, Aluminium-, Granat-Laser (Nd:YAG), mit einer Wellenlänge von 1064 nm, durchgeführt. Über einen Zeitraum von fünf Jahren wurden 82 Patienten behandelt. Eingesetzt wurde der Laser mit 6 W Leistung, Frequenz 40 Hz, 150 nmJ Energie und 100 ms Pulsweite. Der Eingriff wurde in direktem Kontakt mit dem Fettgewebe durch eine Lichtleitfaser in einer 1 mm Kanüle, durchgeführt. Für die Histologie wurden Proben des Fettgewebes sofort nach dem Eingriff entnommen und ein Biopsie vier Wochen nach Eingriff.

Die Submentale laserassistierte Liposuktion ergab ein aussagekräftiges kosmetisches Ergebnis. Die Histologie zeigte eine Ruptur der Adipozyten Membrane mit der daraus folgenden Freisetzung zellulärer Inhalte in den extrazellulären Raum. Collagen-Koagulation sowie separate Kanäle wurden entlang des Pfades der Laserkanüle beobachtet. Zusätzlich wurden kleine Blutgefäße im Fettgewebe koaguliert. Subjektiv analysiert, waren Komplikationen und Ergebnisse ähnlich wie bei traditionellen Lipoplastie-Techniken: eine adäquate Hautkontraktion mit daraus resultierender Verbesserung der Wölbung des zervicofacialen Winkels. Der Laser bewies sich als effektiv bei zellulärer Lysis, Collagenneubildung und bei der Reorganisation der retikulären Dermis. Der Nd:YAG Laser ist ein sehr hilfreiches Gerät für die Behandlung lokaler Lipidistrophie.

Summary

Submental liposuction is an excellent procedure for improving the aging neck by reducing the cervicomental angle and increasing the definition of the mandibular border. Liposuction is currently one of the most common cosmetic surgical procedures. Advances in cannula design, new concepts, and different technical options have improved patient outcomes. Concerns still exist regarding total aspirated volume, and the clinical and hemodynamic risks it entails to the patient, as well as post-operative recovery and outcome. Liposuction of the neck and jowls is clinically one of the most rewarding procedures in aesthetic surgery. In the present study, the author describes his experience of neck and jowl laser-assisted liposuction and its effects on fatty and surrounding tissue.

In this procedure, submental localized lipodystrophy is treated with a neodymium, yttrium, aluminum, garnet (Nd:YAG) laser, at a wavelength of 1064nm. Over a 5-year-period, 82 subjects were treated using 6W power, 40Hz frequency, 150mJ energy and 100ms pulse width parameters. Treatment was performed in direct contact with the fatty tissue via optical fiber delivered through a 1mm cannula. Histology was performed on samples of fatty tissue obtained immediately following the procedure and from biopsy 4 weeks following the procedure.

Submental laser-assisted liposuction resulted in significant cosmetic improvement. Histology revealed rupture of the adipocyte membrane with the consequent release of cellular contents into the extracellular space. Collagen coagulation and discrete channels were observed along the path of the laser cannula. Additionally, small blood vessels were coagulated in the fat tissue. Subjectively analyzed, complications and outcomes were similar to those obtained using traditional lipoplasty techniques. An adequate skin contraction was observed in all treated patients with a consequent improvement in the concavity of the cervicofacial angle. The laser proved to be effective for cellular lysis, collagen neoformation and for the reorganization of the reticular dermis. The direct contact Nd:YAG laser is a useful tool for the treatment of local lipodystrophy.

Introduction

Accumulation of submental fat causes the cervicofacial concavity to decrease, eventually approaching a flat angle or convexity. This may be more apparent as the skin loses elasticity and begins to hang. Submental liposuction is not only a procedure with low-risk but also a relatively inexpensive one. This treatment can offer dramatic changes to the appearance of the neck and mandibular border. With recent technical advances, the results achieved with this surgical treatment are safer and more effective.

Liposuction has become increasingly popular over the last decade and is now among the most popular body sculpting procedures. This increasing popularity is associated with the evolution of techniques and equipment for fat removal and body reshaping. Besides the traditional suction-assisted lipoplasty, other options include ultrasound-assisted and external ultrasound-assisted liposuction, power-assisted liposuction, laser lipolysis as well as low-level laser-assisted liposculpture. The efforts in the search for alternatives and new tools aim mainly at reducing downtime, decreasing operator effort for the surgeon and assistant, reducing bleeding, promoting skin tightening, and facilitating treatment of fibrous or reoperative areas.

Recently, lasers have been adapted for the treatment of localized fat. The laser-fat interaction was initially described by Apfelberg [1–3] in 1992. Publications by Blugerman, Schavelzon and Goldman [4–9] demonstrated their experience with lasers used directly on adipose tissue in a procedure called laserlipolysis. Later, Badin [10, 11] published his experience with laserlipolysis, highlighting the important tissue retraction obtained with this technique. Neira [12] analyzed the probable formation of a pore on the adipocyte membrane after the action of an external laser. However, the result was questioned and non-reproducible in a publication by Brown [13]. In a recent study, Ichikawa [14] showed the histological evaluation in subjects and laboratory animals treated with laserlipolysis.

The mechanism of action of laserlipolysis is selective photothermolysis. In this process, laser light energy is converted into heat energy when absorbed by fat. Conducted by a flexible fiber optic delivered through a cannula, the laser energy is transmitted to the adipocytes, which absorb the energy, expand their volume and rupture. The action time of the laser varies according to the area to be treated and tissue resistance. All subjects suitable for a traditional liposuction method can be treated with laserlipolysis.

Subjects and Methods

A total of 82 consecutive subjects who underwent the submental laserlipolysis procedure between March 1999 and November 2004, at Clínica Goldman de Cirurgia Plástica, in Porto Alegre, Brazil, were included in the study. All underwent pre-operative assessment to determine their general medical condition. None were obese and all were non-smoking and in normal or well controlled health. All subjects provided informed consent. The procedure was recommended for patients with neck lipodystrophy, without strong vertical platysmal bands, and without important skin laxity.

The procedures were performed after subcutaneous infiltration of a solution containing 20 cc of 2% lidocaine with vasoconstrictor and 20 to 30 cc of warm saline solution. The procedure was initiated fol-

Tab. 1: Demographics of Subjects Undergoing Submental Laserlipolysis

Number of Subjects	82	
Average Age (Y)	35.53	(18–61)
Males	4	(5 %)
Females	78	(95 %)

lowing a twenty-minute delay, allowing for diffusion of infiltrate and appropriate vasoconstriction. Of the 82 subjects, 4 (5 %) were men, and 78 (95 %) were women. Ages ranged from 18 to 61 years, with an average of 35.53 years (Tab. 1). All procedures were performed using local anesthesia or local anesthesia assisted by anesthesiologist.

Histological studies were performed in some subjects, focusing on the effects of the laser on the vessels, fat and dermis while using different colorations such as Orcein and Hematoxylin eosin. Biopsies were obtained immediately after laser irradiation and 40 days following the procedure, with the consent of the patient.




Fig. 1: 1 mm cannula containing fiber optic extended approximately 2 mm from the distal end, and emitting laser energy.

Laser

A 1064 nm Nd:YAG laser at 40 Hz and 150 mJ, 6W and 100-microsecond pulse duration was used (Deka, Calenzano, Italy). In this procedure, the laser energy is conducted to the adipose tissue through a 300-micron fiber optic, delivered through a 1 mm diameter stainless steel micro-cannula of variable length. The distal portion of the fiber optic is extended approximately 2 mm beyond the distal end of the cannula (Fig. 1). For visualization, a He:Ne laser source is combined into the beam path. The association of the 1064 nm laser with a He:Ne source allows precise visualization of the region where the energy is acting, due to cutaneous trans-illumination. A 1 mm incision is enough to introduce the cannula, which is then moved in the fat tissue at various depths, including the subdermal layer, similar to a superficial liposuction. When in contact with the previously infiltrated fatty tissue, the light energy produced by the laser is absorbed and converted to heat, thereby expanding the adipocyte contents and rupturing the cell membrane. Thus, through the phenomenon of photothermolysis, cellular rupture is produced. A photoacoustic effect may also play a role in cellular lysis, due to the rapid absorption by and heating of the cell.

The laser treatment is delivered over a varying length of time, and total deposited energy goes according to the size of the treated area, and tissue resistance. In areas of fibrosis or previously treated zones, the treatment time is typically longer. The equipment provides accumulated energy in joules, which can be used as a reference for treatment of similar areas in different subjects. Depending on the volume of treated fat, an oily solution, the product of the cellular lysis, was removed using negative pressure of around 450 mmHg in conjunction with a suction cannula. Aspiration of laserlipolysis debris was only not performed in cases where the treated area contained very small volumes of fat, or in subjects where the goal was to produce superficial relaxation for the purpose of tissue adjustment or cellulite attenuation. Submental laserlipolysis was performed as an outpatient procedure and after the first post-operative day the patients gradually returned to normal daily activities, typically with little discomfort. In some cases, a smooth compression was used for one week. On the second day, the patient initiated treatment with manual lymphatic drainage, external ultrasound, low power external laser and other physiotherapeutic cares. Complications included 2 cases of asymmetry. No case of injury to the marginal mandibular nerve was observed in this series. Subjects were treated in the outpatient setting and were discharged immediately after the procedure or a few hours following the procedure. The careful and judicious selection and preparation of subjects, as well as proper subject orientation, were essential for the success and safety of the procedure.

Histological Studies and Results

The histological assessment was carried out on tissues removed from the subjects immediately following the procedure, and on biopsies taken approximately 40 days following the procedure. Significant findings following laserlipolysis included the coagulation of small blood vessels in the fatty tissue (Fig. 2), the rupture of adipocytes, the appearance of small channels produced by laser action (Fig. 3), reorganization of the reticular dermis (Fig. 4) and coagulation of collagen in the fat tissue (Fig. 5). Although the final result obtained with laserlipolysis was similar to the author's experi-




Fig. 2: Histology showing coagulation of small blood vessels in fat tissue by means of laser action (Orcein 32 x).




Fig. 3: Channel produced by Nd:YAG laser, showing the rupture of adipocytes and fragments of adipose cell membranes (Hematoxylin eosin 32 x).




Fig. 4: Reorganization of reticular dermis. (Hematoxylin eosin 32 x)




Fig. 5: Collagen Coagulation (Hematoxylin eosin 100 x)

ence with traditional liposuction methods, the histological findings suggest several positive benefits brought by the use of the Nd:YAG laser (Fig. 8 to 11), which include skin retraction – due to collagen neoformation – and a reduction in perioperative and postoperative bleeding as well as in the population of adipocytes. Complications were similar to those found with other lipoplasty methods and, in this study, there were no side effects directly related to the laser use.

Discussion

The aim of this study was to analyze the effects of the direct Nd:YAG laser irradiation on fat tissue and the results of this procedure – laserlipolysis in the submental region. There are few studies in the literature showing laser action on lipodystrophy. The techniques published to date using laser described both external and internal methods. This method of lipoplasty exhibited results similar to those from other traditional body sculpting methods previously used by the author. Internal laserlipolysis with 1064 nm Nd:YAG laser has been proving to be a safe and effective method. The decrease in tissue trauma is likely to be associated with the laser-induced coagulation of small vessels in the fat tissue, the adequate infiltration of the

Smartlipo

Interstitial Laser-lipolysis System



Source of Beauty

- Reduced recovery times.
- Minimum post-operative side effects.
- Less invasive and less traumatic technique compared to traditional procedures.
- Safety ensured for both patient and operator.
- Minimum interference with the patient's social life.



DEKA-LMS
DEKA-LMS Laser- & Medizin-Systeme Vertrieb GmbH
 DEKA-LMS Lasersysteme GmbH
 Am Wald 1, 85354 Freising
 Tel. 0 81 67 / 95 00 90
 Fax 0 81 67 / 95 00 99
 stephan.gruner@deka-lms.de

Einführungs-
preise in
Deutschland!




Fig. 6 and 7: A 40-year-old woman with submental lipodystrophy. Preoperative views. (Left). Postoperative views after 7 months (Right).



Fig. 8: A 43-year-old man. Preoperative view. (Left). Postoperative view after 5 months. (Right).



Fig. 9 and 10: A 36-year-old patient who underwent laser-assisted liposuction on the submental region. Preoperative views. (Left). Postoperative views after 3 months. (Right).

anesthetic solution and the possibility of using smaller caliber cannulas. The treatment of areas where fat removal is difficult, as is the case of gynecomastia and previously operated areas, is facilitated by the small diameter of the microcannula containing the laser fiber. The thermal action of the beam facilitates movement of the cannula, consequently entailing less effort and fatigue for the surgeon. The visualization of a reddish color from the He-Ne source, due to its transillumination through the skin, makes the procedure very precise, given that the surgeon knows exactly where the laser is acting. The potential to use the laser for superficial treatment, as proposed by Bolivar [15], may represent another option for the treatment of cellulite, remodeling collagen and smoothing the treated area. Although it still represents a recent technical option, several authors have demonstrated similar results in histological evaluations, describing the effects of the Nd:YAG laser on adipocytes. The main findings are the rupture of the adipocyte cell membrane, the formation of channels, and the coagulation of the collagen. These factors may be responsible for the observed tissue retraction following the procedure. Although some studies have not demonstrated the value of low intensity laser acting transcutaneously, these findings cannot

be applied to the technique described, which employs laser energies far in excess of those employed in low-level techniques. The laser energy from the Nd:YAG used in this study acts directly in contact with the adipose cells, making it unnecessary to cross the skin barrier. Although several authors have corroborated the findings of this study, further clinical studies are still necessary in order to better define the value and limitations of the technique.

Conclusion

The histological findings and postoperative clinical outcomes of subjects who underwent laser lipolysis of the submental area, neck, and jowl have proven the safety and effectiveness of this procedure. Although the laser lipolysis using a Nd:YAG laser has been shown to be effective and safe in this series, further studies are necessary to improve and clarify the utility of this technique.

Acknowledgements

The author thanks Dr. Diego Schavelzon and Dr. Guillermo Blugerman for their invaluable scientific collaboration.



Fig. 11 and 12: A 31-year-old woman with submental lipodystrophy. Preoperative views. (Left). Postoperative views after 7 months. (Right).

Author:

Senior member of the Brazilian Society of Plastic Surgery
Member of the International Society of Aesthetic Plastic Surgery
Member of the Brazilian Society of Laser in Medicine and Surgery

Address of correspondence:

Alberto Goldman, MD
Av. Augusto Meyer 163 conj. 1203 CEP
BR-90550-110 Porto Alegre RS Brazil
alberto@goldman.com.br

Literature

- 1 Apfelberg D (1992) Laser-assisted liposuction may benefit surgeons and subjects. Clin. Laser Mon. 10:259.
- 2 Apfelberg D, and Rosenthal S, Hunstad J (1994) Progress Report on Multicenter Study of Laser-Assisted Liposuction. Aesthetic Plast. Surg. 18(3): 259-64.
- 3 Apfelberg D et al. (1996) Results of Multicentric Study of Laser-Assisted Liposuction. Clin. Plast. Surg. 23(4): 713-9.
- 4 Blugerman G. (2000) Laserlipolysis for the treatment of localized adiposity and "cellulite". Abstracts of World Congress on Liposuction Surgery. Dearborn, Michigan, USA.
- 5 Schavelzon D, Blugerman G, Goldman A, et al. (2001) Laser lipolysis. Abstracts of the 10th International Symposium on Cosmetic Laser Surgery. Las Vegas, USA.
- 6 Goldman A, Schavelzon D, Blugerman G (2002) Laser lipolysis: liposuction using Nd:YAG laser. Revista da Sociedade Brasileira de Cirurgia Plástica. 17: 17-26.
- 7 Goldman A, Schavelzon D, Blugerman G. (2002) Laserlipólise - lipos aspiração com Nd:YAG laser. Revista da Sociedade Brasileira de Laser em Medicina e Cirurgia. 2(5).
- 8 Goldman A (2002) Lipoaspiração a laser - laserlipólise no contorno corporal. Revista Brasileira de Cirurgia. 92.
- 9 Goldman A, Schavelzon D, Blugerman G (2003) Liposuction using neodymium:yttrium-aluminum-garnet laser. Abstract in Plast. Reconstr. Surg. 111:2497.
- 10 Badin A, Moraes L, Godek L, et al. (2002) Laser lipolysis: flaccidity under control. Aesth. Plast. Surg. 26:335-339.
- 11 Badin A, Moraes L, Godek L, et al. (2002) Laserlipólise: flacidez sob controle. Revista da Sociedade Brasileira de Laser em Medicina e Cirurgia.
- 12 Neira R, Arroyave J, Ramirez H, et al. (2002) Fat liquefaction: effect of low-level laser energy on adipose tissue. Plast. Reconstr. Surg. 110:912.
- 13 Brow S, Rohrich R, Kenkel J, et al. (2004) Effect of low-level laser therapy on abdominal adipocytes before lipoplasty procedures. Plast. Reconstr. Surg. 113:1796.
- 14 Ichikawa K, Miyasaka M, Tanaka R, et al. (2005) Histologic evaluation of the pulsed Nd:YAG laser for laser lipolysis. Laser Surg. Med. 36: 43-46.
- 15 De Souza Pinto EB, Erazo PJ, MUNIZ AC et al. (1996) Superficial Liposuction, Aesth. Plast Surg. 20: 111-112.